

**Evaluating the Tea Bag Index approach for different management practices in agroecosystems using long-term field experiments in Austria and Sweden**

Maria Regina Gmach<sup>1\*</sup>, Martin A. Bolinder<sup>1</sup>, Lorenzo Menichetti<sup>1</sup>, Thomas Kätterer<sup>1</sup>, Heide Spiegel<sup>2</sup>, Olle Åkesson<sup>1,3</sup>, Jürgen Kurt Friedel<sup>4</sup>, Andreas Surböck<sup>4</sup>, Agnes Schweinzer<sup>5</sup>, Taru Sandén<sup>2</sup>

<sup>1</sup>Swedish University of Agricultural Science (SLU), Department of Ecology, Box 7044, 75007, Uppsala, Sweden

<sup>2</sup>Austrian Agency for Health & Food Safety (AGES), Department for Soil Health and Plant Nutrition, Spargelfeldstraße 191, A-1220 Vienna, Austria

<sup>3</sup>Lantmännen Lantbruk, Mariestadsvägen 104, 541 39 Skövde, Sweden

<sup>4</sup>University of Natural Resources and Life Sciences (BOKU), Department of Sustainable Agricultural Systems, Institute of Organic Farming (IFÖL), Gregor-Mendel-Straße 33, A-1180 Vienna, Austria

<sup>5</sup>Easy-Cert services GmbH, Königsbrunner Straße 8, Austria

Corresponding author

\* Maria Regina Gmach

Swedish University of Agricultural Science (SLU)

Uppsala, Sweden.

E-mail: gmachmr@gmail.com

Phone: +55 49991164271 / +49 17635956337

## Abstract

Litter decomposition is an important factor affecting local and global C cycles. It is known that decomposition through soil microbial activity in ecosystems is mainly influenced by soil type and climatic conditions. However, for agroecosystems, there remains a need for a better understanding how management practices influence litter decomposition. This study examined the effect of different management practices on decomposition at 29 sites with long-term (mean duration of 38 years) field experiments (LTEs) using the Tea Bag Index (TBI) protocol with standard litter (Rooibos and Green tea) developed by Keuskamp et al. (2013). The objective was to determine if the TBI decomposition rate ( $k$ ) and stabilization factor ( $S$ ) are sensitive enough to detect differences in litter decomposition between management practices, and how they interact with edaphic factors, crop type and local climatic conditions. Tea bags were buried and collected after ~90 days in 16 Austrian and 13 Swedish sites. The treatments at Austrian LTEs focused on mineral and organic fertilizer application, tillage systems and crop residues management, whereas those in Sweden addressed cropping systems, mineral fertilizer application and tillage systems. The results showed that in Austria, incorporation of crop residues and high N fertilizer application increased  $k$ , compared to crop residues removal and low or no N application, respectively. Minimum tillage had significantly higher  $k$  compared to reduced and conventional tillage. In Sweden, fertilized plots showed higher  $S$  than non-fertilized plots and high N fertilizer had the highest  $k$ . Growing spring cereal lead to higher  $k$  than forage crops. Random Forest regressions for Austria and Sweden jointly showed that  $k$  and  $S$  were mainly governed by climatic conditions, which explained more than 70% of their variation. However, under similar climatic conditions, management practices strongly influenced decomposition dynamics. It would be appropriate to apply the TBI approach in a more large-scale network on LTEs for agroecosystems, in order to improve its usefulness as an indicator for the effect of management practices on litter decomposition dynamics, particularly

~~linking it with the potential for C storage as an indicator to better assess, which of the management practices can best promote a higher soil C sink.~~

## **Introduction**

Soil organic carbon (SOC) is one of the most used indicators for soil quality, since its dynamics is involved in regulating ecosystem functionality through its influences on physical, biological and chemical soil properties, which are critical for nutrient cycling and soil fertility (Davidson et al., 2006; Janzen, 2015). Management practices, such as fertilizer application, use of catch- and cover crops, organic amendments, length of bare fallow periods, permanent surface protection with perennial crops, tillage practices and aboveground crop residue management, are impacting SOC balances for agroecosystems (Kätterer and Bolinder, 2022; Sandén et al., 2018; Paustian et al., 2016).

Agricultural soils play a crucial role in the global carbon (C) cycle due to their C sink capacity (Paustian et al., 2016; Lal, 2004). In this context, improved management practices that maintain or increase SOC stocks are considered essential in national greenhouse gas reporting systems (IPCC, 2006), as well as in other international incentives such as the 4 per mille initiative (Minasny et al., 2017). The SOC balance is dynamic and determined by the difference between annual C inputs to soil, and the annual C outputs through the decay of existing soil organic matter and resulting from microbial activity, which is the main contributor to stable SOC (Tiefenbacher et al., 2021; Bolinder et al., 2007). Management practices have a great impact on these two factors by affecting either the amount of C inputs or outputs through decomposition, or both factors simultaneously.

Litter decomposition is a complex biogeochemical process controlled by several biotic and abiotic factors, where the biological activity of decomposers varies with soil properties and is driven largely by climatic conditions (Daebeler et al., 2022; Bradford et al., 2016; Cleveland

et al., 2014; Gholz et al., 2000). Decomposition and SOC stabilization are long-term processes, therefore long-term field experiments (LTEs) are among the most useful resources for quantifying the impact of management practices on litter decomposition, SOC changes, and soil functioning (Sandén et al., 2018; Kätterer et al., 2012; Bergkvist and Öborn, 2011). Within LTE's, experiments determining litter mass loss over time in situ are important for understanding SOC dynamics, nutrient cycling and colonization by soil biota under field conditions. The traditional method that has been used in ecology for more than 50 years consists of litterbag studies, burying known quantities of various organic materials into the soil, and retrieving them successively at different intervals (Kampichler and Bruckner, 2009; Burgess et al., 2002; Bock and Gilbert, 1957). These studies are not always comparable because they are subject to variations in e.g., litter type, mesh-size, sample preparation and analytical methods, and the placement of litterbags may alter the microclimate for decomposers (Kampichler and Bruckner, 2009).

Keuskamp et al. (2013) developed therefore a standardized, low-cost and time-efficient methodology called Tea Bag Index (TBI), characterizing the decomposition process with commercially available tea bags, where green tea is representing labile organic material and rooibos tea as a surrogate for recalcitrant litter. A decomposition rate ( $k$ ) and a stabilization factor ( $S$ ) are obtained accordingly with their chemical composition and the respective weight lost at a single point in time after an incubation period of ca 90-days in the soil. The TBI approach is particularly useful for assessing geographical differences in decomposition dynamics because results are directly comparable across sites, varying only with local edaphic and seasonal environmental conditions (Keuskamp et al., 2013).

In several studies, the TBI has been used as an indicator for biological (Sandén et al., 2021; Costantini et al., 2018) and microbial activity (Daebeler et al., 2022; Treharne et al., 2019; Tóth et al., 2017), ecosystem functioning, nutrient cycling (Zaller et al., 2016), and soil quality

(Tresch et al., 2018; Buchholz et al., 2017). Generally, this follows the concepts of soil quality as reviewed by Bünemann et al. (2018), where TBI would primarily be a biological soil quality indicator. According to the TBI community, data collected from networks of researchers and citizen scientists for constructing a global TBI map ([www.teatime4science.org](http://www.teatime4science.org)) shows that most studies have been using the TBI approach for different forest and grassland ecosystems or urban soils, studies in agricultural fields represent less than 15% of the TBI database. Indeed, there are only a few published studies (Daebeler et al., 2022; Dossou-Yovo et al., 2022; Struijk et al., 2022; Fu et al., 2021; Sandén et al., 2020; Barel et al., 2019; Poeplau et al., 2018; Sievers and Cook, 2018) that have been using the TBI approach for evaluating agroecosystems, and it is not clear if this method is sensitive enough to detect differences between management practices.

This study used the TBI approach to investigate the effect of management practices on the decomposition rate ( $k$ ) and stabilization factor ( $S$ ) at several LTEs in Austria and Sweden, with different soil characteristics, and climatic conditions, and subjected to various management practices.

To the best of our knowledge, this is the first analysis using the TBI approach for such a large number of LTEs and management practices for agroecosystems. The management practices included organic amendments, crop rotations, aboveground crop residue handling, mineral fertilizer application, and tillage. Our objectives were: (i) to evaluate if the TBI  $k$  and  $S$  parameters are sensitive enough to distinguish litter decomposition between different management practices; (ii) to quantify the effect of management practices on  $k$  and  $S$ ; and (iii) to identify the most important local climate and/or soil properties affecting litter decomposition in Austria and Sweden.

## Materials and Methods

## Study sites

### *Austria*

We used sixteen Austrian (AT) sites, by selecting contrasting management practices (as treatments) from three different categories of LTEs where the management practices had been in place for 11 to 63 years (see details in Table 1). The TBI measurements were made in 2014, 2015 and 2016. Measurements sometimes took place in more than one year at the same LTE (e.g., MUBIL), and the sites were abbreviated as AT1 to AT16. Six experiment categories involved carbon balance practices (CB) focusing on organic matter inputs such as compost and crop residues, eight sites were studying soil fertility (SF) in terms of differences in mineral N and P fertilization, and two sites examined tillage systems (TS). The sites are located in several agricultural areas across the country (Fig. 1), with diverse soil textures (Table S1) and variable crop types (Table 1) and climatic characteristics (Table 3) during the years of TBI measurements. More details for some of the sites are available in specific publications: AT3 (Spiegel et al., 2018; Lehtinen et al., 2017; Tatzber et al., 2015; Aichberger and Söllinger, 2009), AT4 to AT6 (Spiegel et al., 2018; Lehtinen et al., 2014), AT15 and AT16 (Tatzber et al., 2015; Spiegel et al., 2007). In addition, Sandén et al. (2018) puts the Austrian LTEs in the context of other European LTEs.

The purpose of the Austrian C balance LTEs was threefold: i) to assess the sustainability of stockless vs. livestock-keeping organic farming management on soil and crop traits (AT1, AT2), ii) to investigate the effects of compost amendments on soil and crops (AT3), and iii) to compare crop residue incorporation and removal (AT4-AT6). Compost amendment LTE and crop residue incorporation LTEs also included mineral fertilizer application, whereas AT1 and AT2 only focused on different organic fertilizer treatments.

The eight soil fertility LTEs (AT7-AT14) were all focusing on the effect of mineral fertilizer on soil and crop properties. In most cases, treatments studied different amounts of mineral nitrogen fertilizer, whereas AT9 and AT12 also investigated the effect of different amounts of K fertilizer application. Nitrogen fertilizer was applied in four stages and potassium in three stages, according to Austrian guidelines for fertilizer (BMLFUW, 2017).

For the tillage system experiments, conventional tillage (CT) was compared to reduced tillage (RT) and minimum tillage (MT). Regular mouldboard ploughing to 25–30 cm soil depth was applied in CT treatment, whereas cultivator in autumn to a depth of 15–20 cm was used in RT treatment and a rotary driller that loosened the soil to a depth of 5–8 cm was used in MT treatment. The soil was turned over only in the CT treatment, where inversion tillage was incorporating the crop residues. Fertilizer application was crop specific according to the Austrian guidelines for fertilizing (BMLFUW, 2017).

## *Sweden*

We used thirteen Swedish (SE) sites by selecting contrasting treatments from three different categories of LTEs, where the management practices had been in place for 11 to 59 years (see details in Table 2). The TBI measurements at these sites were made only in one year (2016) and were abbreviated as SE1 to SE13. Six sites involving combined management practices (CMP), four studying the effect of rotations (ROT), and three sites with tillage systems (TS). The sites are located in several agricultural areas across the country (Fig. 1), with diverse soil textures and variable crop types (Table 2) and climatic characteristics (Table 3) during the year of TBI measurements. Bergkvist and Öborn (2011) give a general description of all these LTEs. Only a brief description of treatments are provided below, more details on the sites with combined management practices is given by Carlgren and Mattson (2001), and for tillage

systems by Arvidsson et al. (2014), while Poeplau et al. (2015) provide some more insight on the rotation experiments.

The initial purpose of the six LTEs with combined management practices (SE1-SE6) was to compare a change from the traditional mixed farm production system including crops and livestock into a pure cash crop system, by studying their effects on the sustainability of crop production and soil properties (entitled soil fertility experiments). The dairy production treatments contain exclusively perennial grass-clover leys and receive one farmyard manure (FYM) application per rotation. The cash crop treatments consist of annual crops (i.e., oilseed is replacing leys in the rotation) without manure applications (0 FYM) only receiving mineral fertilizers application (NPK). The PK applications in all treatments we selected were aimed at achieving rapid build-up of the soil PK status, i.e., the amount applied was first replacing that exported in harvested products (i.e., maintenance principle), to which an extra amount was added (corresponding to the max treatment). The N-rates in all NPK treatments were also corresponding to max application rate, and were adapted depending on crop type, where spring cereals, oilseeds, and leys received 125 kg, while sugar beet received 210 kg N ha<sup>-1</sup> yr<sup>-1</sup>. We were also using the control plots receiving no NPK (0 NPK). As a third factor, aboveground crop residue removal takes place in all FYM treatments, simulating use of harvest residues for fodder or bedding material that are recycled as manure. The southern sites have 4-year rotations and those in central Sweden have 6-year rotations. The north site (SE5) is slightly different from the others, consisting of a 7-year rotation and is studying only the livestock-based production system.

For rotation experiments purposes, we were comparing extreme treatments representing two rotations from four LTEs (SE7-SE10) with the main objective to study changes in SOC (named humus balance experiments), i.e., a continuous spring cereal (SC) system and a ley-dominated rotations (L). The straw was removed from the plots every year in the SC treatments, and L



consisted of a grass-clover mixture re-established every fourth year. Both rotations were receiving P and K accordingly with the maintenance principle, and SC and L were receiving 120 and 150 kg N ha<sup>-1</sup> yr<sup>-1</sup>, respectively.

In the three tillage experiments (SE11-SE13), the conventional tillage (CT) and direct seeding (DS) treatments were the same for all sites, consisting of inversion plowing to a depth of 20-23 cm and by using a disc seed drill, respectively. The shallow (5-7 cm) and deep (~12 cm) reduced tillage treatments (SRT and DRT, respectively), consisted of primary tillage operations made in the autumn and most commonly with a chisel plough. The main crops in all the tillage system experiments were winter and spring cereals (occasionally oilseed), fertilized accordingly with local recommendations and with the aboveground residues chopped and left in the field.

#### **TBI method and sampling design**

The TBI method was used according to the protocol established by Keuskamp et al. (2013) to determine litter decomposition using two types of commercial tetrahedron-shaped tea bags by Lipton Unilever (Green tea and Rooibos tea). The green tea (*Camellia sinensis*; EAN: 8722700055525) has high cellulose content, higher soluble fraction, and lower C:N ratio; while rooibos tea (*Aspalathus linearis*; EAN: 8722700188438) has high lignin content, lower soluble fraction, and higher C:N ratio, which is expected to slow down decomposition (Keuskamp et al., 2013). The synthetic tea bag material has a mesh size opening of 0.25 mm allowing access to microorganisms, very fine roots and root hairs.

The initial mass of the tea bag contents was determined on 20 randomly selected bags for each tea type from different boxes, oven-dried at 70°C for 48 hours and weighed separately; the mean dry mass for green tea was  $1.717 \pm 0.048$  g and that for rooibos tea was  $1.835 \pm 0.027$  g.

222 For both countries, close to seeding of annual crops, from end of April to mid-June depending  
223 on location, each tea bag was properly identified and buried in the soil at 8 cm depth.

224 For Austria, four bags of each tea were used and placed side by side at a distance of 2 to 3 cm.  
225 Each tea bag was properly identified and buried in the soil at 8 cm depth. The TBI incubation  
226 period from placement to last retrieval averaged  $80 \pm 13$  days (Table 3) due to logistic issues for  
227 collection. After collecting, the tea bags were oven-dried at  $70^\circ\text{C}$  for 48 hours after removal of  
228 adhered soil particles according to the standardized protocol by Keuskamp et al. (2013). After  
229 drying, the tea bags were opened, and the tea content was weighted. The ash content was not  
230 determined.

231 The same TBI protocol was used for the Swedish sites. As in Austria, four bags of each tea  
232 were used per experimental unit, placed side-by-side at a distance of 2 to 3 cm. Each tea bag  
233 was properly identified and buried in the soil at 8 cm depth. The mean TBI incubation period  
234 from placement to last retrieval date averaged  $91 \pm 1$  day (Table 3). In addition to the removal  
235 of adhering soil particles, the ash content was determined (i.e., both for green and rooibos tea  
236 on mixed samples of the four replicates) in a muffle oven at  $550^\circ\text{C}$  for 16 hours. The rationale  
237 for measuring the ash content was that three of the Swedish sites had a high clay content (Table  
238 S1), where the complete removal of adhering soil particles may be more difficult. However,  
239 the ash content was on average quite low, representing  $15 \pm 6\%$  and  $10 \pm 4\%$  for the Green and  
240 Rooibos tea bags, respectively (data not shown).

241 After measuring the remaining dry matter, the decomposition rate ( $k$ ) and stabilization factor  
242 ( $S$ ) for both countries were calculated according to the TBI presented by Keuskamp et al.  
243 (2013). This standardized method that is using single measurements after an incubation period  
244 in the soil of 90-days have received some criticism. For instance, Mori (2022) and Mori et al.  
245 (2023) showed that this incubation period is not always long enough for the mass loss of green  
246 tea to reach a plateau, and further suggested that time-series mass loss data of rooibos tea is

also required to respect the underlying assumptions of the TBI method. Time-series (15, 30, 60 and 90-days) of green and rooibos tea were available for all the Swedish sites but only at one Austrian site (16, 26, 62 and 91-days at AT16). The incubation period was consistently always 90-days for the Swedish sites, and only shorter than that (i.e., about 60-days incubation period) for a few of the Austrian sites (Table 3). To have as uniform comparisons as possible between the two datasets, we only used the last measurement for both countries for calculating  $k$  and  $S$ . The purpose of using the time-series for testing the underlying TBI assumptions was beyond the scope of this paper.

The daily climate data for Austria were retrieved from the Central Institution for Meteorology and Geodynamics (ZAMG). For Sweden, the daily climate data were gathered through official data from the most nearby LantMet climate stations, and from the Swedish Meteorological and Hydrological Institute (SMHI). The climate variables used in this study were air temperature, precipitation, solar radiation, wind speed and air humidity (Table 3).

For Austrian soils, pH was measured electrochemically (pH/mV Pocket Meter pH 340i, WTW, Weilheim, Germany) in 0.01 M CaCl<sub>2</sub> at a soil-to-solution ratio of 1:5 (ÖNORM L1083). Total soil organic C (TOC) concentrations were analyzed by dry combustion in a LECO RC-612 TruMac CN (LECO Corp., St. Joseph, MI, United States) at 650°C (ÖNORM L1080). Total N (N<sub>tot</sub>) was determined according to ÖNORM L1095 with elemental analysis using a CNS (carbon, nitrogen, sulfur) 2000 SGA-410–06 at 1250°C. Texture was determined according to ÖNORM L1061-1 and L1062-2. For Sweden the data were gathered from recent archived analysis protocols (pH was measured in water). Clay content, C content, C:N ratio, and pH measured from each site in both countries are shown in Table S1 (Supplementary material).

## **Data analysis**

271 Analysis of variance (ANOVA) for each experiment category (i.e., CB, SF, TS, CMP and ROT)  
272 was performed to analyze the effects of the treatments and the differences between sites on  $k$   
273 and  $S$  separately for both countries. When the treatments were identical within the same  
274 experiment category, sites were used as a random effect with a mixed ANOVA to test the  
275 average treatment effect, mean values were used as replicates to test the differences between  
276 sites. The Tukey's test ( $p < 0.05$ ) was used for comparing the same treatments and the same  
277 sites using R software version 4.2.2. We have treated the data from different years at the same  
278 sites in Austria as independent observations, in the sense they are not a time series of  
279 measurement. Interactions between site and treatment were considered.

280 We calculated a climate-dependent soil biological activity parameter ( $Re_{clim}$ ), by using mean  
281 daily air temperature, total precipitation, and potential evapotranspiration (PET) data in  
282 pedotransfer, soil water balance and biological activity functions. Compared to raw climatic  
283 data alone, this parameter is integrating the effect of climate, soil and crop properties. It is  
284 calculated as the product of a soil temperature ( $Re_{temp}$ ) and relative water content ( $Re_{wat}$ ) factor  
285 with a daily time step (i.e.,  $Re_{clim} = Re_{temp} \times Re_{wat}$ ), which is thereafter averaged to give an  
286 estimate of soil biological activity for a given time period. These two factors are derived from  
287 soil temperature and soil moisture response functions expressing the activity of decomposers  
288 and their relative effect on the decay rates of organic materials in the arable layer of agricultural  
289 soils. Briefly, the  $Re_{temp}$  is calculated from air temperature and leaf area index using an  
290 empirical model (Kätterer and Andrén, 2009), while  $Re_{wat}$  is calculated using pedotransfer  
291 functions for simulating the soil water balance and a function for estimating PET. In addition  
292 to air temperature and leaf area index, calculations of  $Re_{wat}$  also involve the use of daily climatic  
293 data for precipitation, wind speed, air humidity, and solar radiation, as well crop types and  
294 yields, soil texture and SOC content. For details see Bolinder et al., 2008, and Fortin et al.,  
295 2011) and information in the following R package used for the calculations:

296 <https://github.com/ilmenichetti/reclim> (Please refer to the package documentation). The *Re<sub>clim</sub>*  
297 concept can be used for quantifying regional differences in soil biological activity alone  
298 (Bolinder et al., 2013; Andrén et al. 2007) or integrated as a parameter in the ICBM SOC model.  
299 In the latter case, it is adjusting the decomposition rates of both C inputs to soil from crop  
300 residues (e.g., straw) and that of the more stable SOC (Andrén and Kätterer, 1997; Andrén et  
301 al., 2004). In this study, we used the concept of *Re<sub>clim</sub>* and the *Re<sub>temp</sub>* and *Re<sub>wat</sub>* factors to test if  
302 the product of soil-temperature and relative water content better explained the variation in *k*  
303 and *S* than the two latter alone, and to determine if they also better explained this variation  
304 compared to using only raw climatic data.

305 We calculated simple correlation between variables mean annual temperature (MAT), mean  
306 temperature during the incubation period (MAT<sub>TBI</sub>), total annual precipitation (TAP), total  
307 precipitation during incubation period (TP<sub>TBI</sub>), potential evapotranspiration (PET), potential  
308 evapotranspiration during incubation period (PET<sub>TBI</sub>), aridity index (AI), aridity index during  
309 incubation period (AI<sub>TBI</sub>), temperature and precipitation factor (TxP), *Re<sub>clim</sub>*, *Re<sub>wat</sub>*, *Re<sub>temp</sub>*, pH,  
310 SOC, clay and C:N ratio, using Pearson correlation. For more accurate results, we applied  
311 random forest (RF) regression in order to rank the importance of variables for *k* and *S*, using  
312 the random forest R package (Liaw and Wiener, 2002). The RF is a machine learning technique  
313 based on decision trees that predicts a certain variable from a set of other variables through a  
314 series of binary splits of the data, where the variables are either continuous or categorical. For  
315 example, in the case of a continuous variable it consists of all the data points above or below a  
316 certain threshold, for a categorical variable it consists of all the data points belonging or not to  
317 a specific class. All these subsequent splits constitute a decision tree. A random forest is a set  
318 of decision trees and it is therefore an ensemble technique. This allowed us to utilize treatment  
319 and crop variables (including N fertilizer) without having to convert them into a ranking.  
320 Another useful asset of an RF regression is that it evaluates the importance of each variable in

defining the predicted variable. There are various possible measurements to do that, but they are all based on measuring the effectiveness of each subsequent split in each node of a decision tree in sorting out the information. In our study, we used a measurement called node purity based on the Gini index, which expresses the probability of one split of the data (i.e., one node of the tree) defining the predicted variable. The total node purity of a certain variable in a tree is the sum of all the node purity measurements for each node considering that particular variable, and the higher it is the more that variable is important.

We used the following models to predict the two TBI parameters  $k$  and  $S$ :

$$k_n f(TN, SOC, CN, cl, cr, tr, pH, N, PET, PET_{TBI}, TAP, TP_{TBI}, MAT, MAT_{TBI}, AI, AI_{TBI}, Re_{clim}, Re_{wat}, Re_{temp})$$

$$S_n f(TN, SOC, CN, cl, cr, tr, pH, N, PET, PET_{TBI}, TAP, TP_{TBI}, MAT, MAT_{TBI}, AI, AI_{TBI}, Re_{clim}, Re_{wat}, Re_{temp})$$

where soil variables (continuous) were TN (total N g kg<sup>-1</sup>), SOC (g kg<sup>-1</sup>), CN (C:N ratio), cl (clay content g 100 g<sup>-1</sup>) and pH. Categorical variables were N (N fertilizer application factor with 4 levels), cr (crop factor, e.g., barley, ley (establishment), ley (production), oat, spring oilseeds, sugar beet, maize and winter wheat) and tr (a treatment factor with 30 levels). The climatic variables (PET<sub>TBI</sub>, PET, TP<sub>TBI</sub>, TAP, MAT<sub>TBI</sub>, MAT, AI<sub>TBI</sub> and AI) are as defined in Table 3. The climate response variables  $Re_{clim}$ ,  $Re_{wat}$ ,  $Re_{temp}$  are as described above.

Since many of variables in our model are likely to be correlated and carry similar information, we applied the recursive feature elimination algorithm implemented in the caret R package by Kuhn et al. (2016), which assess in subsequent iterations the optimal set of predicting variables (features) to be utilized by the RF model. The procedure starts by fitting a RF model with all variables, ranking them by importance, and discarding the least important. The algorithm then iterates. The optimal number and set of features are then defined by a fitness metric (in our case the model R<sup>2</sup>), selecting the set with the best model fitness. The selected models were used to compute the variables' relative importance.

In order to better understand the similarities between the sites, we run a principal component analysis (PCA) in the space composed by the variables pH, SOC, TN, C:N, clay, TAP, MAT, PET, AI and TxP. The analysis was run with the R command `prcomp` (Venables and Ripley, 2002), from the base R installation (“Stats” package). Variables were all standardized by rescaling them to zero mean and unit variance.

## Results

### *Effect of management practices*

#### *Austria*

Both the TBI parameters  $k$  and  $S$  varied between treatments and sites in Austria, and even between years at the same site within the C balance category (Fig. 2 and Table S2). In general, all treatments in AT1 with a lucerne crop under wetter conditions (2014) showed higher  $k$  and lower  $S$  than in AT2 with a wheat crop under dryer conditions (2015), and the treatment with municipal compost and green manure (FW) had the highest  $S$  in 2015. The AT3 site did not show significant differences between the treatments. Crop residue incorporation treatment (CRI) had higher  $k$  than the crop residue removal (CRR) at the AT4 and AT6 sites, and AT6 showed a higher  $k$  than at AT4. Comparing years for the same experiment site and type with different crops, AT5 (2015; with wheat) had higher  $S$  than AT6 (2016; with maize).

For the soil fertility experiment category (Table 1), AT12 had the highest  $k$  and AT13 had the highest  $S$ . Sites receiving NPK fertilizer application (AT7, AT8 and AT9) had higher  $k$  and  $S$  even at different N doses than at the AT12, AT13 and AT14 sites receiving only N (i.e., without P and K). Stabilization was significantly higher in AT9 than in AT7. For potassium trials, AT11 showed significantly higher  $k$  and  $S$  than AT10. Regarding sites receiving only N addition (AT12, AT13, and AT14), maximum doses (180 kg N) showed the highest  $k$  (0.0095), and no N addition had the lowest  $k$  (0.0066) (Fig. 2 and Table S2).

Regarding the tillage system experiment category (AT15 and AT16 - Table 1),  $k$  was significantly higher in shallow reduced tillage (SRT) and  $S$  was higher in deep reduced tillage (DRT) in 2015 (AT15), but no significant differences between treatments were found in 2016 (AT16). Site AT16 had significantly higher  $S$  than AT15 (Fig. 2. and Table S2).

#### *Sweden*

At the Swedish sites (Table 2), the TBI measurements for the combined management practices experiment category showed that both  $k$  and  $S$  were significantly higher for the FYM/NPK treatments maximum amount of farmyard manure and maximum doses of NPK (FYM/NPK) (Fig. 3, Table S3) compared with the control treatments (0 FYM/0 NPK). Comparing sites, SE5 and SE6 had the highest  $k$  and SE4 had the lowest  $k$ , while SE3 showed the highest  $S$  followed by SE6, SE5 and SE4. The lowest  $S$  was for SE1 and SE2 (Table S3).

Regarding the rotation experiments, the continuous spring cereal rotation showed higher  $k$  than for ley, but there was no significant difference in  $S$ . Comparing sites, SE9 showed higher  $k$  than SE7, SE8 and SE10, whereas SE7 and SE8 had the highest  $S$ .

For the tillage system experiment category, conventional tillage (CT) had the lowest  $k$  and  $S$ , while deep reduced tillage (DRT) had the highest  $k$  and  $S$ . The highest  $S$  was observed for the SE12 followed by SE11 and SE13. Sites did not show significant differences for  $k$ .

Comparing tillage system experiments in Austria and Sweden (2016) for sites (AT16, SE11, SE12 and SE13) for the treatments that were the same in both countries (CT, SRT and DRT), deep reduced tillage had the highest  $k$  and shallow reduced tillage had the highest  $S$ . The Austrian site showed the lowest  $k$  and the highest  $S$  compared to the Swedish sites, which did not show significant differences among them (Table S3).

The field application of the TBI found a clear discrimination of both  $k$  and  $S$  values between the two countries. The both mean  $k$  and  $S$  values were higher in Sweden (Table 4, Fig. 4). In



general, the variation in  $k$  and  $S$  values were lower in Austria. Indeed, mean  $k$  by site in Austria varied between 0.0058 and 0.0128 and mean  $S$  varied between 0.113 and 0.442 (Table S2). Whereas mean  $k$  by site in Sweden was between 0.0084 and 0.0301, and mean  $S$  was between 0.118 and 0.361 (Table S3). All values for  $k$  and  $S$  were within the range of the previous global TBI investigation (0.005-0.04 for  $k$ ; and 0.05-0.55 for  $S$ ) by Sandén et al. (2020).

### *Influence of climate and soil properties*

Using the combined dataset for Austria and Sweden, resulted in significant negative correlation between  $k$  and MAT, TAP, PET, TxP factor,  $Re_{clim}$  and  $Re_{temp}$ , and significant positive correlation with the C:N ratio (Table S4). Stabilization factors in Austria and Sweden combined correlated negatively with MAT<sub>TBI</sub> period,  $Re_{clim}$  and  $Re_{temp}$ .

The variable selection procedure with the random forest models (Fig. 5) identified fewer variables explaining  $k$  and  $S$  values for the combined dataset. The variable explaining  $k$  is low, but the overall predicting power of the model decreased substantially.

According to the optimized random forest model, most of the variance in  $k$  was accounted for by climatic variables (Fig. 5), with  $Re_{temp}$  ranking the highest, followed by AI and MAT. It was also showing that treatment, C:N ratio and SOC were important factors as well. The  $S$  was also influenced by climate-related variables, with  $Re_{temp}$  leading the ranking, and it was followed by pH and more agronomic variables such as treatment, crop and N fertilizer application.

The principal component analysis (Fig. 6) revealed that the data are well divided in two groups, representing the two countries, which allow us to say that indeed edapho-climatic characteristics are different between Austria and Sweden. The PC1 showed that points in Austria are positively related to climatic characteristics, such as MAT, TAP and PET. On the other hand, points in Sweden are positively influenced by high clay content and high C:N ratio. High C:N enhancement is related to low TAP and TxP. High clay contents are related to low

MAT and PET, which can be due to an historical influence on soil weathering. The PC2 was dominated by SOC and TN contents, mainly at two points in Austria (AT7 and AT9: two sites testing soil fertility with NPK and relatively low TAP and  $TP_{TBI}$  compared to other sites). Austria showed more divergent data, having more heterogeneous sites than in Sweden.

## Discussion

### *Influence of climate and soil properties*

In previous studies using the TBI approach, it was shown that climate played a significant role on decomposition in a temperate biome (Djukic et al., 2018), but when comparing several different biomes, climatic conditions were of relatively low importance (Fanin et al., 2020). In Boreal soils, Althuizen et al. (2018) found that increased temperatures enhanced  $k$ , whereas increased precipitation decreased  $k$  across years. Many studies have shown a positive correlation between precipitation and decomposition rates (Pimentel et al., 2019; García Palacios et al., 2016), however precipitation did not show correlation with  $k$  according to the random forest analysis. On the other hand,  $Re_{wat}$  showed great importance (Fig. 5). It is because this variable includes nonlinearities due to its shape according to which decomposition increases with soil moisture and then decreases at high soil water content due to oxygen limitation of microorganisms (Moyano et al., 2013).

In general, higher  $k$  values were observed when the aridity index was lower, and it was identified by the random forest regression model being an important variable affecting the rate of decomposition (Fig. 5). This may be related to the observations that in more arid and warmer sites, the biological processes driving SOC dynamics are impaired which may result in decreasing soil C and N stocks (Kerr and Ochsner, 2020; Ontl and Schulte, 2012; Jiao et al., 2016; Reynolds et al., 2007).

For practical reasons the teabags in our study were buried in the soil during the growing season, corresponding to a period when the soil biological activity is highest (Bolinder et al., 2013). When burying the teabags during the growing season, the difference in climatic variables between sites are attenuated, in particular with respect to air temperature. For example, the MAT at the Swedish most northerly (SE5 and SE9) and southerly (SE2) sites are 4.1 and 9.0 °C, respectively, whereas the corresponding mean air temperatures during the incubation period (MAT<sub>TBI</sub>) were 14.3 and 16.2 °C, respectively.

The TBI  $S$  parameter was also dependent on climate. Indeed, the random forest model identified climatic parameters as the main factors affecting  $S$  in both Austria and Sweden during all evaluated periods. In particular,  $Re_{clim}$  and  $Re_{temp}$  often showed significant negative correlations, which implies a negative impact of air temperature on  $S$ . However, raw climatic variables, such as precipitation and temperature were only weakly correlated with  $S$ . This is probably due to nonlinear processes, which are considered in the ICBM climate-dependent soil biological activity calculations such as  $Re_{wat}$  (as discussed above). Furthermore, since litter decomposition dynamics is influenced by multiple factors that interact and change over time (Bradford et al., 2016), the relationships are not always linear. Random forest models that we fitted to the data are more efficient capturing such combinations and interactions of factors, and can detect relationships that would not be detectable by linear approaches.

#### *Effect of management practices*

Our results revealed that a large number of management practices significantly affected both the decomposition rate  $k$  and stabilization factor  $S$  according to the TBI approach used in several LTEs in Austria and Sweden (Fig. 2, 3 and 5). This is in contrast to the studies by Djukic et al. (2018) as well as Saint-Laurent and Arsenault-Boucher (2020), who did not find any

significant effect of land use and management practices on early-stage litter decomposition in a temperate biome.

In the C balance trials in Austria, soils receiving green manure + municipal compost had higher  $S$  than soil receiving biogas slurry at the second year (Table S2). This is in agreement with studies indicating that compost can improve SOC stabilization over time (Mekki et al., 2019; Eshetu et al., 2013; Ceccanti et al., 2007). The higher  $k$  in the crop residues incorporated treatments (AT4 and AT6; Table S2) can be attributed to the fact that incorporation of crop residues into the soil can increase the decomposition rate by stimulating microbial activity. During the early stages of decomposition, soluble C is rapidly utilized by soil biota (Werth and Kuzyakov, 2010). The higher  $k$  and lower  $S$  at AT6 compared to AT4 were likely due to the loamy texture, lower potential evapotranspiration resulting in a lower aridity index at the AT4 site (Table 3 and Table S1), corroborated by the PCA analysis (Fig. 6).

There were no significant differences in  $k$  and  $S$  found among treatments in the soil fertility trials in Austria with NPK addition. However, there was a trend towards a higher  $S$  at AT9 compared to AT7, likely related to the higher SOC content in AT9, since the climatic conditions and soil texture were quite similar for both areas, which suggests that higher SOC content may have increased  $S$ . Site AT11 had higher  $k$  and  $S$  than AT10. Possible explanations for this trend are that AT10 had lower clay content, lower precipitation resulting in higher PET (corroborated by the PCA analysis) and AI, contributing to a lower soil moisture content and thereby lower decomposition and stabilization.

Nitrogen supply may favor microbial activity and thereby litter decomposition (Raiesi, 2004). This was reflected in the treatments where only N was added, where the high dose of 180 kg N ha<sup>-1</sup> (AT12, AT13 and AT14) induced a significantly higher  $k$  (Table S2), compared to the treatments with no N addition, which also had the lowest  $k$ . Furthermore, the significant

difference between sites, in which AT12 had the highest  $k$  could at least partly be explained by a higher SOC and higher pH at this site (corroborated by the PCA analysis).

In the tillage system experiment at Fuchsenbigl in Austria in 2015 (AT15, with maize) and 2016 (AT16; with wheat), the shallow reduced tillage showed significantly higher  $k$  than deep reduced tillage and conventional tillage, only in 2015, indicating that shallow soil tillage stimulated decomposition that particular year. This was likely due to climatic conditions, since 2015 was slightly drier and warmer, furthermore maize straw has lower C:N ratio than wheat straw and tends to decompose faster. Some studies showed faster decomposition under conventional tillage than under reduced tillage practices (e.g., Lupwayi et al., 2004). However, Kainiemi et al. (2015) found a decrease in soil respiration in conventional tillage compared to shallow tillage in temperate regions, which directly implies a lower decomposition (and lower  $k$ ). These differences between tillage treatments are attributable to indirect effects on soil moisture and temperature profiles. We attribute the significantly higher  $S$  in 2016 to the fact that this year was moister and less warm, compared to 2015, resulting in a lower aridity index during the TBI incubation period ( $AI_{TBI}$ ).

In the Swedish combined management practices trials, treatments receiving organic and mineral fertilizer application had higher  $k$  and  $S$  (FYM/NPK; Table S3), likely due to the increase in microbial diversity and activity favored by nutrient and C supply (Stark et al., 2007). Sites SE5 and SE6 showed the highest  $k$ : SE5 had low evapotranspiration and aridity index, resulting in more moisture; SE6 had also high  $S$ , due to high clay content and low PET (corroborated by the PCA analysis). Site SE3 had high  $S$ , which could be related to a higher C:N ratio, as suggested in another TBI experiment by Althuizen et al. (2018), a study showing that C:N ratio is positively correlated to  $S$ . The SE1 and SE2 sites had lower  $S$  than SE3 despite similar climatic conditions, which probably was related to the type of crops growing in these

treatments (i.e., sugar beet in the SE1 and SE2 and grass/clover ley in SE3), since these two crop types have different effects on soil temperature and moisture.

In the Swedish rotation system trials, spring cereal (SC) had higher  $k$  than ley (Table S3). Site SE9 had higher  $k$  and lower  $S$ , in which the low stabilization may be caused by low clay content, low pH, and high solar radiation, leading to low SOC (corroborated the PCA analysis). The highest  $S$  were found in sites SE7 and SE8, in which the former showed high clay and SOC content, and SE8 had high precipitation and low PET.

For the tillage system treatments in Sweden, similar to the Austrian sites, conventional tillage showed the lowest  $k$ , and also lowest  $S$ . Even when comparing tillage systems in Sweden and Austria jointly (Table S3) we noticed that conventional tillage also showed the lowest  $k$ , while deep reduced tillage the highest.

As expected, the decomposition rate  $k$  was mostly affected by climatic conditions such as temperature and aridity or moisture. However agronomic and edaphic factors (i.e., soil management choice (treatment), soil C:N ratio, SOC, crop type, pH, clay content, soil nitrogen and others)- proved to be of great relevance for  $k$ . The stabilization factor  $S$  expresses the degree by which the labile fraction of the plant material is decomposed. We observed that the influence of pH was the most important edaphic factor, and the agronomic factors as the soil management treatment, crop type, and N fertilizer application were also identified as good predictors. A study conducted by Fu et al. (2021) suggested that pH, nutrient availability and soil compaction were the main reasons contributing to the differences in litter decomposition. The net effect of pH is not clear since it modifies both SOC decay kinetics and productivity simultaneously (Paradelo et al., 2015), with a maximum effect at around neutral pH (Liao et al., 2016). Nevertheless, the impact of pH on litter decomposition using the TBI approach seems clear in our study.

## Conclusion

Our results show that both the TBI  $k$  and  $S$  parameters were sensitive to management practices in agroecosystems in Austria and Sweden. We observed significant differences for some of the treatments in all categories of LTEs. Notably, for the effect of crop residue incorporation, organic amendments and N fertilizer application, crop types and tillage systems. In the Austrian LTEs, application of green manure + municipal compost showed a higher  $S$  compared to the application of other organic amendments. Incorporation of crop residues instead of crop residue removal and high N fertilizer instead of low or no N application increased decomposition rate  $k$ . In the Swedish LTEs, it was shown that combined management practices with both farmyard manure and mineral NPK resulted in higher  $k$  and  $S$  compared to no manure and no NPK applications, whereas growing spring cereals instead of leys increased  $k$  but did not change  $S$ . For both countries, tillage systems with deep reduced tillage practices had higher  $k$ , and shallow reduced tillage had higher  $S$ . However, these effects were also site or year dependent within a given country. Climatic conditions had the most important impact on the decomposition rate  $k$  and the stabilization factor  $S$ , but also pH, treatment, crop types, SOC, C:N ratio and clay content were good predictors of the TBI parameters. Generally, the correlations with raw climatic variables such as precipitation and temperature were quite poor. Better relationships were found when nonlinearities due to interactions between climatic and edaphic conditions were accounted for. The results highlight how a wide range of management practices used in agroecosystems affect TBI parameters jointly to soil and climatic conditions. This also implies suggests that the TBI  $k$  and  $S$  parameters could serve as indicators of how different agricultural management practices influence the global carbon cycle via decomposition, a matter requiring further in-depth investigation., ~~and is useful to assess their potential contribution to act as a sink for atmospheric C.~~

568    **Data availability**

569    Data can be provided by the authors upon request.

570

571    **Author contribution**

572    MRG, MB and TS wrote and prepared the manuscript draft; MB, TK and TS supervised and

573    led the research; MB, OA, HS, JKF, AS, AS and TS developed the methodology; MRG and

574    LM analyzed the data. All authors reviewed and edited the manuscript.

575

576    **Competing interests**

577    The contact author has declared that none of the authors has any competing interests.

578

579    **Acknowledgements**

580    Financial support was provided by the Swedish Farmers' Foundation for Agricultural Research,

581    grant number O-18-23-141. Part of this research has been done in the framework of the EJP

582    SOIL that has received funding from the European Union's Horizon 2020 research and

583    innovation programme: Grant agreement No 862695.

584



## 585    **References**

- 586    Aichberger, K., Söllinger, J.: Use of biocompost in agriculture – results of a long-term field  
587    trial. In: Spiegel, H, Zonno, V, editors. Realising the ETAP in the management of waste from  
588    farms, proceedings of the second AQUAGRIS workshop Vienna, 19th June 2009. Vienna:  
589    AGES; p. 6–8., 2009.
- 590    Althuizen, I. H. J., Lee, H., Sarneel, J. M., Vandvik, V.: Long-term climate regime modulates  
591    the impact of short-term climate variability on decomposition in Alpine grassland soils.  
592    *Ecosystems*, 21, 1580-1592, DOI:10.1007/s10021-018-0241-5, 2018.
- 593    Andrén, O., Kätterer, T.: ICBM: The introductory carbon balance model for exploration of soil  
594    carbon balances, *Ecol Appl*, 7, 1226-1236, DOI: 10.1890/1051-0761, 1997.
- 595    Andrén, O., Kätterer, T., Karlsson, T.: ICBM regional model for estimations of dynamics of  
596    agricultural soil carbon pools, *Nutr Cycl Agroecosyst* 70, 231-239, DOI:  
597    10.1023/B:FRES.0000048471.59164.ff, 2004.
- 598    Andrén, O., Kihara, J., Bationo, A., Vanlauwe, B., Kätterer, T.: Soil climate and decomposer  
599    activity in Sub-Saharan Afrika estimated from standard weather station data: A simple climate  
600    index for soil carbon balance calculations, *Ambio*, 36, 379-386, DOI: 10.1579/0044-  
601    7447(2007)36[379:scadai]2.0.co;2, 2007.
- 602    Arvidsson, J., Håkansson, I.: Response of different crops to soil compaction—Short-term  
603    effects in Swedish field experiments, *Soil Till Res*, 138, 56-63, DOI:  
604    10.1016/j.still.2013.12.006, 2014.
- 605    Barel, J. M., Kuyper, T. W., Paul, J., de Boer, W., Cornelissen, J. H. C., De Dein, G. B.: Winter  
606    cover crop legacy effects on litter decomposition act through litter quality and microbial  
607    community changes, *J Appl Ecol*, 56, 132-143, DOI:10.1111/1365-2664.13261, 2019.
- 608    Bergkvist, G., Öborn, I.: Long-term field experiments in Sweden—what are they designed to  
609    study and what could they be used for, *Aspects of Applied Biology*, 113, 75-85, 2011.
- 610    BMLFUW: Richtlinien für die sachgerechte Düngung. Bundesministerium für Land- und  
611    Forstwirtschaft, Umwelt und Wasserwirtschaft, 2017.
- 612    Bockock, K. L., Gilbert, O. J. W.: The disappearance of leaf litter under different woodland  
613    conditions, *Plant Soil*, 9, 179-185, DOI: 10.1007/BF01398924, 1957
- 614    Bolinder, M. A., Andrén, O., Kätterer, T., Parent, L. E.: Soil organic carbon sequestration  
615    potential for Canadian Agricultural Ecoregions calculated using the Introductory Carbon  
616    Balance Model, *Can J Soil Sci*, 88, 451-460, DOI: 10.4141/CJSS07093, 2008.
- 617    Bolinder, M. A., Fortin, J. G., Anctil, F., Andrén, O., Kätterer, T., de Jong, R., Parent, L. E.:  
618    Spatial and temporal variability of soil biological activity in the Province of Québec, Canada  
619    (45-58 oN, 1960-2009) – calculations based on climate records. *Climatic Change*, 117, 739-  
620    755, DOI: 10.1007/s10584-012-0602-6, 2013.
- 621    Bolinder, M. A., Janzen, H. H., Gregorich, E. G., Angers, D. A., VandenBygaart, A. J.: An  
622    approach for estimating net primary productivity and annual carbon inputs to soil for common

623 agricultural crops in Canada, *Agr Ecosyst Environ*, 118, 29–42, DOI:  
624 10.1016/j.agee.2006.05.013, 2007.

625 Bradford, M. A., Berg, B., Maynard, D. S., Wieder, W. R., Wood, S. A.: Understanding the  
626 dominant controls on litter decomposition. *J Ecol*, 104, 229–38, DOI: 10.1111/1365-  
627 2745.12507, 2016.

628 Buchholz, J., Querner, P., Paredes, D., Bauer, T., Strauss, P., Guernion, M., Scimia, J., et al.:  
629 Soil biota in vineyards are more influenced by plants and soil quality than by tillage intensity  
630 or the surrounding landscape, *Sci Rep*, 7, 17445, DOI: 10.1038/s41598-017-17601-w, 2017.

631 Bünemann, E. K., Bongiorno, G., Bai, Z., Creamer, R., de Deyn, G. B., et al.: Soil quality – A  
632 critical review, *Soil Biol Biochem* 120, 105-125, DOI: 10.1016/j.soilbio.2018.01.030, 2018.

633 Burgess, M. S., Mehuys, G. R., Madramootoo, C. A.: Decomposition of grain-corn residues  
634 (*Zea mays* L.): A litterbag study under three tillage systems. *Can J Soil Sci*, 82, 127-38. DOI:  
635 10.4141/S01-013, 2002.

636 Carlgren, K., Mattsson, L.: Swedish soil fertility experiments, *Acta Agric Scandin* 51, 49-76.  
637 DOI: 10.1080/090647101753483787, 2001.

638 Ceccanti, B., Masciandaro, G., Macci, C.: Pyrolysis-gas chromatography to evaluate the  
639 organic matter quality of a mulched soil, *Soil Till Res*, 97, 71-8, DOI:  
640 10.1016/j.still.2007.08.011, 2007.

641 Cleveland, C. C., Reed, S. C., Keller, A. B., Nemergut, D.R., O'Neill, S. P., Ostertag, R.,  
642 Vitousek, P.M.: Litter quality versus soil microbial community controls over decomposition: a  
643 quantitative analysis, *Oecologia* 174, 283-294, DOI: 10.1007/s00442-013-2758-9, 2014.

644 Costantini, E. A. C., Castaldini, M., Diago, M. P.: Effects of soil erosion on agroecosystem  
645 services and soil functions: A multidisciplinary study in nineteen organically farmed European  
646 and Turkish vineyards, *J Environ Manag*, 223, 614-624, DOI: 10.1016/j.envman.2018.06.065,  
647 2018.

648 Daebeler, A., Petrová, E., Kinz, E., Grausenburger, S., Berthold, H., Sandén, T., Angel, R.:  
649 Pairing litter decomposition with microbial community structures using the Tea Bag Index  
650 (TBI), *Soil*, 8, 163-76, DOI: 10.5194/soil-8-163-2022, 2022.

651 Davidson, E. A., Janssens, I. A., Luo, Y.: On the variability of respiration in terrestrial  
652 ecosystems: moving beyond Q<sub>10</sub>, *Glob Change Biol*, 12, 154-64, DOI:10.1111/j.1365-  
653 2486.2005.01065.x, 2006.

654 Djukic, I., Kopfer-Rojas, S., Schmidt, I. K., Larsen, K. S., et al.: Early stage litter  
655 decomposition across biomes, *Sci Total Environ*, 628-629, 1369-1394, DOI:  
656 10.1016/j.scitotenv.2018.01.012, 2018.

657 Dossou-Yovo, W., Parent, S. E., Ziadi, N., Parent, E., Parent, L. E.: Tea Bag Index to assess  
658 carbon decomposition rate in cranberry agroecosystems, *Soil Syst*, 5, 44, DOI:  
659 10.3390/soilsystems5030044, 2022.

660 Eshetu, B., Baum, C., Leinweber, P.: Compost of different stability affects the molecular  
661 composition and mineralization of soil organic matter, *Open J Soil Sci*, 3, 58-69. DOI:  
662 10.4236/ojss.2013.31007, 2013.

663 Fanin, N., Bezaud, S., Sarneel, J. M., Cecchini, S., Nicolas, M., Augusto, L.: Relative  
664 importance of climate, soil and plant functional traits during the early decomposition stage of  
665 standardized litter, *Ecosystems*, 23, 1004-1018, DOI: 10.1007/s10021-019-00452-z, 2020.

666 Fortin, J. G., Bolinder, M. A., Anctil, F., Kätterer, T., Andrén, O., Parent, L. E.: Effects of  
667 climatic data low-pass filtering on the ICBM temperature- and moisture-based soil biological  
668 activity factors in a cool and humid temperate climate, *Ecol Model*, 222, 3050-3060. DOI:  
669 10.1016/j.ecolmodel.2011.06.011, 2011.

670 Fu, Y., Jonge, L. W., Greve, M. H., Arthur, E., Moldrup, P., Norgaard, T., Paradelo, M.: Linking  
671 litter decomposition to soil physicochemical properties, gas transport, and land use, *Soil*  
672 *physics and hydrology - Soil Sci Soc Am J*, 86, 34-46, DOI: 10.1002/saj2.20356, 2021.

673 García Palacios, P., Shaw, E. A., Wall, D. H., Hättenschwiler, S.: Temporal dynamics of biotic  
674 and abiotic drivers of litter decomposition, *Ecol Lett*, 19, 554–563, DOI: 10.1111/ele.12590,  
675 2016.

676 Gholz, H. L., Wedin, D. A., Smitherman, S. M., Harmon, M. E., Parton, W. J.: Long-term  
677 dynamics of pine and hardwood litter in contrasting environments: toward a global model of  
678 decomposition, *Glob Change Biol*, 6, 751–765, DOI: 10.1046/j.1365-2486.2000.00349.x,  
679 2000.

680 IPCC. Agriculture, forestry and other land use. IPCC guidelines for national greenhouse gas  
681 inventories. In: Eggleston, S., Buendia, L., Miwa, K., Ngara, T., Tanebe, K. Institute for Global  
682 Environmental Strategies, prepared by the National Greenhouse Gas Inventories Programme,  
683 Hayama, Kanagawa, Japan, 2006.

684 Janzen, H. H.: Beyond carbon sequestration: soil as conduit of solar energy, *Eur J Soil Sci*, 66,  
685 19-32, DOI: 10.1111/ejss.12194, 2015.

686 Jiao, F., Shi, X. R., Han, F. P., Yuan Z. Y.: Increasing aridity, temperature and soil pH induce  
687 soil C-N-P imbalance in grasslands, *Sci Rep*, 6, 19601, DOI:10.1038/srep19601, 2016.

688 Kainiemi, V., Arvidsson, J., Kätterer, T.: Effects of autumn tillage and residue management on  
689 soil respiration in a long-term field experiment in Sweden, *J. Plant Nutr. Soil Sci.*, 178, 189-  
690 198, DOI: 10.1002/jpln.201400080, 2015.

691 Kampichler, C., Bruckner, A.: The role of microarthropods in terrestrial decomposition: a meta-  
692 analysis of 40 years of litterbag studies, *Biol Rev*, 84, 375-89, DOI: 10.1111/j.1469-  
693 185X.2009.00078.x, 2009.

694 Kätterer, T., Andrén, O.: Predicting daily soil temperature profiles in arable soils in cold  
695 temperate regions from air temperature and leaf area index, *Acta Agr Scand*, 59, 77-86. DOI:  
696 10.1080/09064710801920321, 2009.

697 Kätterer, T., Bolinder, M. A.: Chapter 15: Agriculture practices to improve soil carbon  
698 sequestration in upland soil, In: *Understanding and fostering soil carbon sequestration* (ed. Dr  
699 Cornelia Rumpel), DOI: 10.19103/AS.2022.0106.15, 2022.

700 Kätterer, T., Bolinder, M. A., Berglund, K., Kirchmann, H.J.: Strategies for carbon  
701 sequestration in agricultural soils in northern Europe, *Acta Agr Scand*, 62, 181-98. DOI:  
702 10.1080/09064702.2013.779316, 2012.

703 Kerr, D. D., Ochsner, T. E.: Soil organic carbon more strongly related to soil moisture than soil  
704 temperature in temperate grassland, *Soil Sci Soc Am J*, 84, 587-596, DOI: 10.1002/saj2.20018,  
705 2020.

706 Keuskamp, J. A., Dingemans, B. J., Lehtinen, T., Sarneel, J. M., Hefting, M. M.: Tea bag index:  
707 a novel approach to collect uniform decomposition data across ecosystems, *Methods Ecol Evol*,  
708 4, 1070–1075, DOI: 10.1111/2041-210X.12097, 2013.

709 Kuhn, Max. Contributions from Jed Wing, Steve Weston, Andre Williams, Chris Keefer, Allan  
710 Engelhardt, Tony Cooper, Zachary Mayer, Brenton, Kenkel, the R Core Team, Michael  
711 Benesty, Reynald Lescarbeau, Andrew Ziem, Luca Scrucca, Yuan Tang and Can Candan, *Caret*:  
712 Classification and Regression Training. R package version 6.0-71. [https://CRAN.R-](https://CRAN.R-project.org/package=caret)  
713 [project.org/package=caret](https://CRAN.R-project.org/package=caret), 2016.

714 Lal, R.: Soil carbon sequestration impacts on global climate change and food security. *Science*,  
715 304, 1623–1627, DOI: 10.1126/science.1097396, 2004.

716 Lehtinen, T., Dersch, G., Söllinger, J., Baumgarten, A., Schlatter, N., Aichberger, K., Spiegel,  
717 H.: Long-term amendment of four different compost types on a loamy silt Cambisol: impact  
718 on soil organic matter, nutrients and yields, *Arch of Agronomy and Soil Sci*, 63, 663-673, DOI:  
719 10.1080/03650340.2016.1235264, 2017.

720 Lehtinen, T., Schlatter, N., Baumgarten, A., Bechini, L., Krüger, J., Grignani, C., Zavattaro, L.,  
721 Costamagna, C., Spiegel, H.: Effect of crop residue incorporation on soil organic carbon and  
722 greenhouse gas emissions in European agricultural soils, *Soil Use Manage*, 30, 524-538, DOI:  
723 10.1111/sum.12151, 2014.

724 Liao, K., Wu, S., Zhu, Q.: Can Soil pH Be Used to Help Explain Soil Organic Carbon Stocks?,  
725 *Clean Soil Air Water*, 44, 1685–1689, DOI: 10.1002/clen.201600229, 2016.

726 Liaw, A., Wiener, M.: “Classification and Regression by randomForest.” *R News*, 2, 18-22,  
727 [https://CRAN.R-project.org/doc/Rnews/Liaw and Wiener](https://CRAN.R-project.org/doc/Rnews/Liaw%20and%20Wiener), 2002

728 Lupwayi, N. Z., Clayton, G. W., O’Donovan, J. T., Harker, K. N., Turkington, T. K., Rice, W.  
729 A.: Decomposition of crop residues under conventional and zero tillage, *C J Soil Sci*, 84, 403–  
730 410, DOI: 10.4141/S03-082, 2004.

731 Mekki, A., Aloui, F., Sayadi, S.: Influence of biowaste compost amendment on soil organic  
732 carbon storage under arid climate, *Japca J Air Waste Ma*, 69, 867-877, DOI:  
733 10.1080/10962247.2017.1374311, 2019.

734 Minasny, B., Malone, B. P., McBratney, A. B., et al.: Soil carbon 4 per mille, *Geoderma*, 292,  
735 59-86, DOI: 10.1016/j.geoderma.2017.01.002, 2017.

736 Mori T.: Validation of the Tea Bag Index as a standard approach for assessing organic matter  
737 decomposition: A laboratory incubation experiment. *Ecological Indicators*, 141, 109077, DOI:  
738 10.1016/j.ecolind.2022.109077, 2022.

739 Mori, T., Ono, K., Sakai, Y.: Testing the Tea Bag Index as a potential indicator for assessing  
740 litter decomposition in aquatic ecosystems. *Ecological Indicators*, 152, 110358, DOI:  
741 10.1016/j.ecolind.2023.110358, 2023.

742 Moyano, F. E., Manzoni, S., Chenu, C.: Responses of soil heterotrophic respiration to moisture  
743 availability: An exploration of processes and models, *Soil Biol. Biochem*, 59, 72–85, DOI:  
744 10.1016/j.soilbio.2013.01.002, 2013.

745 Ontl, T. A., Schulte, L. A.: Soil carbon storage, *Nature Education Knowledge*, 3, 2012.

746 Paradelo, R., Virto, I., Chenu, C.: Net effect of liming on soil organic carbon stocks: A review.  
747 *Agr Ecosyst Environ*, 202, 98–107, DOI: 10.1016/j.agee.2015.01.005, 2015.

748 Paustian, K., Lehmann, J., Ogle, S., et al.: Climate-smart soils, *Nature*, 532, 49-57, DOI:  
749 10.1038/nature17174, 2016.

750 Pimentel, L. G., Cherubin, M. R., Oliveira, D. M., Cerri, C. E., Cerri, C. C.: Decomposition of  
751 sugarcane straw: Basis for management decisions for bioenergy production, *Biomass Bioenerg*,  
752 122, 133-44, DOI: 10.1016/j.biombioe.2019.01.027, 2019.

753 Poeplau, C., Kätterer, T., Bolinder, M. A., Börjesson, G., Berti, A., Lugato, E.: Low  
754 stabilization of aboveground crop residue carbon in sandy soils of Swedish long-term  
755 experiments, *Geoderma*, 237, 246-255, DOI: 10.1016/j.geoderma.2014.09.010, 2015.

756 Poeplau, C., Zopf, D., Greiner, B., et al.: Why does mineral fertilization increase soil carbon  
757 stocks in temperate grasslands?, *Agr Ecosyst Environ*, 265, 144-155, DOI:  
758 10.1016/j.agee.2018.06.003, 2018.

759 Raiesi, F.: Soil properties and N application effects on microbial activities in two winter wheat  
760 cropping systems, *Biol Fert Soils*, 40, 88-92, DOI: DOI 10.1007/s00374-004-0741-7, 2004.

761 Reynolds, J. F., Smith, D. M. S., Lambin, E. F., et al.: Global desertification: Building a science  
762 for dryland development, *Science*, 316, 847–851, DOI: 10.1126/science.1131634, 2007.

763 Saint-Laurent, D., Arsenault-Boucher, L.: Soil properties and rate of organic matter  
764 decomposition in riparian woodlands using the TBI protocol, *Geoderma*, 358, 113976, DOI:  
765 10.1016/j.geoderma.2019.113976, 2020.

766 Sandén, T., Spiegel H., Stüger, H. P., Schlatter, N., Haslmayr, H. P., Zavattaro, L., Grignani, C.,  
767 Bechini, L., D'Hose, T., Molendijk, L., Pecio, A., Jarosz, Z., Guzmán, G., Vanderlinden, K.,

768 Giráldez, J. V., Mallast, J., ten Berge, H.: European long-term field experiments: knowledge  
769 gained about alternative management practices. *Soil Use Manage*, 34, 167-176, DOI:  
770 10.1111/sum.12421, 2018.

771 Sandén, T., Spiegel, H., Wenng, H., Schwarz, M., Sarneel, J. M.: Learning science during  
772 teatime: Using a citizen science approach to collect data on litter decomposition in Sweden and  
773 Austria, *Sustainability*, 12, 29-39, DOI:10.3390/su12187745, 2020.

774 Sandén, T., Wawra, A., Berthold, H., Miloczki, J., Schweinzer, A., Gschmeidler, B., Spiegel,  
775 H., Debeljak, M., Trajanov, A.: TeaTime4Schools: Using Data Mining Techniques to Model

776 Litter Decomposition in Austrian Urban School Soils, *Front Ecol Evolut*, 9, 432, DOI:  
777 10.3389/fevo.2021.703794, 2021.

778 Sievers, T., Cook, R. L.: Aboveground and root decomposition of cereal rye and hairy vetch  
779 cover crops, *Soil Sci Soc Am J*, 82, 147-155, DOI: 10.2136/sssaj2017.05.0139, 2018.

780 Spiegel, H., Dersch, G., Hösch, J., Baumgarten, A.: Tillage effects on soil organic carbon and  
781 nutrient availability in a long-term field experiment in Austria, *Die Bodenkultur*, 58, 1-4, 2007.

782 Spiegel, H., Mosleitner, T., Sandén, T., Zaller, J. G.: Effects of two decades of organic and  
783 mineral fertilization of arable crops on earthworms and standardized litter decomposition, *Die*  
784 *Bodenkultur: Journal of Land Management, Food and Environment*, 69, 17-28, DOI:  
785 10.2478/boku-2018-0003, 2018.

786 Stark, C., Condon, L. M., Stewart, A., Di, H. J., O'Callaghan, M.: Influence of organic and  
787 mineral amendments on microbial soil properties and processes, *Appl Soil Ecol*, 35, 79-93,  
788 DOI: 10.1016/j.apsoil.2006.05.001, 2007.

789 Struijk, M., Whitmore, A. P., Mortimer, S., Shu, X., Sizmur, T.: Absence of a home-field  
790 advantage within a short-rotation arable cropping system, *Plant Soil*, 26, 1-7, DOI:  
791 10.1007/s11104-022-05419-z, 2022.

792 Tatzber, M., Schlatter, N., Baumgarten, A., Dersch, G., Körner, R., Lehtinen, T., Unger, G.,  
793 Mifek, E., Spiegel, H.: KMnO<sub>4</sub> determination of active carbon for laboratory routines: three  
794 long-term field experiments in Austria, *Soil Res*, 53, 190-204, DOI: 10.1071/SR14200, 2015.

795 Tiefenbacher, A., Sandén, T., Haslmayr, H-P., Miloczki, J., Wenzel, W., Spiegel, H.: Optimizing  
796 carbon sequestration in croplands: a synthesis, *Agronomy*, 11, 882 DOI:  
797 doi:10.3390/agronomy11050882, 2021.

798 Tóth, Z., Táncsics, A., Kriszt, B., et al.: Extreme effects of drought on decomposition of the  
799 soil bacterial community and decomposition of plant tissue, *Eur J Soil Sci*, 68, 504-513, DOI:  
800 10.1111/ejss.12429, 2017.

801 Treharne, R., Bjerke, J. W., Tømmervik, H., et al.: Arctic browning: Impacts of extreme  
802 climatic events on heathland ecosystem CO<sub>2</sub> fluxes, *Glob Chang Biol*, 25, 489-503, DOI:  
803 10.1111/gcb.14500, 2019.

804 Tresch, S., Moretti, M., Le-Bayon, R. C., Mäder, P., Zanetta, A., Frey, D., et al.: Urban soil  
805 quality assessment – A comprehensive case study dataset of urban garden soils, *Front Environ*  
806 *Sci*, 6, 136, DOI: 10.3389/fenvs.2018.00136, 2018.

807 Venables, W. N., and B. D. Ripley: *Modern Applied Statistics with S*, Springer-Verlag (2002).

808 Werth, M., Kuzyakov, Y.: <sup>13</sup>C Fractionation at the Root-Microorganisms-Soil Interface: A  
809 Review and Outlook for Partitioning Studies, *Soil Biol Biochem*, 42, 1372-1384, DOI:  
810 10.1016/j.soilbio.2010.04.009, 2010.

811 Zaller, J. G., König, N., Tiefenbacher, A., Muraoka, Y., Querner, P., Ratzenböck, A.,  
812 Bonkowski, M., Koller, R.: Pesticide seed dressings can affect the activity of various soil  
813 organisms and reduce decomposition of plant material, *BMC Ecol*, 16, 37, DOI:  
814 10.1186/s12898-016-0092-x, 2016.



## Tables

Table 1. Experimental sites in Austria. Site acronym, location name, year of TBI measurements, category and type of LTE including its duration, main crop cultivated during the TBI measurements and management treatments

Site	Location	Year	Category*	Experiment <sup>†</sup>	Age	Crop	Treatments <sup>§</sup>
AT1	MUBIL	2014	CB	OA	11	Luzerne	GM, FW, FYM, BS
AT2	MUBIL	2015	CB	OA	12	Wheat	GM, FW, FYM, BS
AT3	Ritzlhof	2015	CB	IF-N & OA+N	24	Maize	0, 40, 90, 120 N CFW, CGM, CS, CSS+80N
AT4	Rottenhaus	2016	CB	CR with NPK	31	Maize	CRR, CRI
AT5	Rutzendorf	2015	CB	CR with NPK	34	Wheat	CRR, CRI
AT6	Rutzendorf	2016	CB	CR with NPK	35	Maize	CRR, CRI
AT7	Breitstetten	2015	SF	IF-N with PK	40	Wheat	60, 90, 120, 145 N
AT8	Breitstetten	2016	SF	IF-N with PK	41	Wheat	60, 90, 120, 145 N
AT9	Haringsee	2015	SF	IF-N with PK	40	Wheat	60, 90, 120, 145 N
AT10	Fuchsenbigl	2016	SF	IF-K with NP	61	Maize	0, 150, 300 K
AT11	Rottenhaus	2016	SF	IF-K with NP	63	Maize	0, 150, 300 K
AT12	Haringsee	2016	SF	IF-N without PK	41	Maize	0, 60, 120, 180 N
AT13	Zinsenhof	2016	SF	IF-N without PK	41	Maize	0, 60, 120, 180 N
AT14	Zissersdorf	2016	SF	IF-N without PK	41	Maize	0, 60, 120, 180 N
AT15	Fuchsenbigl	2015	TS	TS	27	Maize	CT, SRT, DRT
AT16	Fuchsenbigl	2016	TS	TS	28	Wheat	CT, SRT, DRT

\* CB: carbon balance practices; SF: soil fertility; TS: tillage systems.

<sup>†</sup> OA: organic amendments; IF-N & OA+N: N inorganic fertilization and organic amendments plus mineral N; CR & IF-P with NK: crop residues and P inorganic fertilization with 90 and 40 kg ha<sup>-1</sup> of N and K, respectively; IF-N with PK: N inorganic fertilization with 55 and 180 kg ha<sup>-1</sup> of P and K, respectively; IF-N without PK: N inorganic fertilization; IF-K with NP: K inorganic fertilization with 40 and 120 kg ha<sup>-1</sup> of P and K; TS: tillage system.

<sup>§</sup> GM: green manure; FW: municipal compost and green manure; FYM: farmyard manure; BS: biogas slurry; CFW: compost food waste with 175 kg N ha<sup>-1</sup>; CGM: compost green manure with 175 kg N ha<sup>-1</sup>; CS: compost slurry with 175 kg N ha<sup>-1</sup>; CSS: compost sewage sludge with 175 kg N ha<sup>-1</sup>; CRR: crop residues removed; CRI: crop residues incorporated; CT: conventional tillage; SRT: shallow reduced tillage; DRT: deep reduced tillage.



Table 2. Experimental sites in Sweden. Site acronym, location name, year of TBI measurements conducted in 2016, type of LTE including its duration, main crop cultivated during the TBI measurements and management treatments

Site	Name	Experiment <sup>*</sup>	Age	Crop	Treatments <sup>†</sup>
SE1	Börgeby	CMP	59	Sugar beet	FYM/NPK
				Sugar beet	0 FYM/0 NPK
SE2	Ekebo	CMP	59	Sugar beet	FYM/NPK
				Sugar beet	0 FYM/0 NPK
SE3	Högåsa	CMP	50	Ley production year	FYM/NPK
				Ley production year	FYM/0 NPK
				Spring oilseed	0 FYM/NPK
				Spring oilseed	0 FYM/0 NPK
SE4	Kungsängen	CMP	53	Oat	FYM/NPK
				Oat	0 FYM/0 NPK
SE5	Röbacksdalen	CMP	47	Barley	FYM/NPK
				Barley	FYM/0 NPK
SE6	Vreta Kloster	CMP	50	Ley production year	FYM/NPK
				Ley production year	FYM/0 NPK
				Spring oilseed	0 FYM/NPK
				Spring oilseed	0 FYM/0 NPK
SE7	Lanna	ROT	35	Oat	SC
				Ley production year	L
SE8	Lönnstorp	ROT	36	Barley	SC
				Ley establish. year	L
SE9	Röbacksdalen	ROT	36	Barley	SC
				Ley establish. year	L
SE10	Säby	ROT	46	Wheat	SC
				Ley establish. year	L
SE11	Lanna	TS	34	Winter wheat	CT, DS
SE12	Säby	TS	11	Barley	CT, SRT, DRT, DS
SE13	Ultuna	TS	19	Barley	CT, DRT

<sup>\*</sup> CMP: combined management practices; ROT: rotation systems; TS: tillage systems.

<sup>†</sup> FYM/NPK: maximum amount of farmyard manure and maximum doses of NPK; 0 FYM/0 NPK: no manure and no NPK application; FYM/0 NPK: maximum amount of farmyard manure and no NPK application; 0 FYM/NPK: no manure and maximum doses of NPK; SC: spring cereal; L: ley; CT: conventional tillage; DS: direct seeding; SRT: shallow reduced tillage; DRT: deep reduced tillage.

Table 3. Annual climatic characteristics for the Austrian and Swedish sites during the entire year of measurements and only during the TBI period (days) corresponding to the period between the date of placement and the last retrieval date of the tea bags.

Site	TBI period	TAP	TP <sub>TBI</sub>	MAT	MAT <sub>TBI</sub>	PET	PET <sub>TBI</sub>	AI	AI <sub>TBI</sub>
	days	mm		°C		mm			
<i>Austria</i>									
AT1	99	756	294	12.5	20.3	811.4	414.0	1.07	1.4
AT2	61	516	65	12.6	18.8	605.1	141.3	1.17	2.2
AT3	80	622	161	12.2	20.0	682.9	295.1	1.09	1.8
AT4 & 11	84	939	384	11.2	20.5	763.4	306.1	0.81	0.8
AT5	63	516	82	12.6	19.4	605.1	148.6	1.17	1.8
AT6	96	735	226	12.0	21.6	854.2	394.0	1.16	1.7
AT7	59	516	81	12.6	19.3	605.1	137.4	1.17	1.7
AT8	84	735	240	12.0	17.1	854.2	334.2	1.16	1.4
AT9	60	516	81	12.6	19.2	605.1	139.6	1.17	1.7
AT10 & 16	92	735	205	12.0	21.4	851.2	378.9	1.15	1.8
AT12	87	735	203	12.0	21.8	854.2	367.4	1.16	1.8
AT13	84	939	327	11.2	20.5	763.4	304.3	0.81	0.9
AT14	82	735	154	12.0	21.3	854.2	319.6	1.16	2.1
AT15	81	516	103	12.6	22.5	605.1	212.9	1.17	2.1
<i>Sweden</i>									
SE1	92	436	141	9.2	16.7	444.6	235.1	1.02	1.7
SE2	91	681	270	9.0	16.5	578.5	297.8	0.85	1.1
SE3	91	458	142	7.9	13.3	618.0	306.4	1.35	2.1
SE4	93	429	126	7.1	15.3	640.0	367.0	1.48	2.9
SE5 & 9	92	526	164	4.1	14.3	396.2	218.2	0.75	1.3
SE6	91	433	158	7.9	13.3	617.5	301.8	1.16	1.9
SE7 & 11	91	412	115	7.4	12.7	514.2	276.4	1.3	2.4
SE8	90	610	224	9.4	14.4	500.2	265.7	0.82	1.2
SE10 & 12	91	429	126	7.1	15.3	358.5	155.7	0.84	1.2
SE13	92	429	126	7.1	15.3	639.1	377.8	1.5	3.0

TAP: total annual precipitation; TP<sub>TBI</sub>: total precipitation during TBI period; MAT: mean annual temperature; MAT<sub>TBI</sub>: mean temperature during TBI period; PET: total annual potential evapotranspiration; PET<sub>TBI</sub>: potential evapotranspiration during TBI period; AI: annual aridity index (PET divided by TAP); AI<sub>TBI</sub>: aridity index during TBI period.

856 Table 4 – Mean values of decomposition rate ( $k$ ) and stabilization factor ( $S$ ) for the TBI  
857 approach after the incubation period.

858


	Mean TBI parameters	
	$k$	$S$
<i>Sweden</i>	$0.0160 \pm 0.01$	$0.247 \pm 0.14$
<i>Austria</i>	$0.0115 \pm 0.004$	$0.228 \pm 0.11$


859

860

## Figures

FIGURE 1






868

869

870 **FIGURE 3**


871



872


873

874 **FIGURE 4**




875

876





880 **FIGURE 6**



881

882

## Figure caption

Figure 1 - Location and environmental zone of the Austrian and Swedish sites.

Figure 2 - Average decomposition rate ( $k$ ) and stabilization factor ( $S$ ) for each treatment and site in Austria. The extents of the box indicate 25<sup>th</sup> and 75<sup>th</sup> percentiles, and the lines represent the 50<sup>th</sup> percentile. Whiskers represent the 10<sup>th</sup> and 90<sup>th</sup> percentiles and outliers are given as open symbols. Green boxes: Carbon balance (CB) experiment; yellow boxes: Soil fertility (SF) experiment; purple boxes: tillage systems (TS) experiment. Site AT1 shows results from 2014. Sites AT2, AT3, AT5, AT7, AT9, and AT15 show results from 2015. And sites AT4, AT6, AT8, AT10, AT11, AT12, AT13, AT14, and AT16 show results from 2016.

Figure 3 - Average decomposition rate ( $k$ ) and stabilization factor ( $S$ ) for each site and treatment in Sweden. The extents of the box indicate 25<sup>th</sup> and 75<sup>th</sup> percentiles, and the lines represent the 50<sup>th</sup> percentile. Whiskers represent the 10<sup>th</sup> and 90<sup>th</sup> percentiles. Green boxes: combined management practices (CMP) experiment; yellow boxes: rotation (ROT) experiment; purple boxes: tillage systems (TS) experiment.

Figure 4 - Distribution of the mean decomposition rate constant ( $k$ ) and the stabilization factor ( $S$ ) for each site in Austria and Sweden. Errors bars represent standard deviation.

Figure 5 - a and b) Variables selection procedure to identify the optimal number of variables to explain the variance of  $k$  and  $S$  with a Random Forest model. The blue point represents the optimal model. c and d) Relative importance of the variables used by each optimized Random

907 Forest model to predict the variance in the  $k$  and  $S$  parameters in Austria and Sweden jointly.  
908 The higher the Node purity, the higher the importance of such variable.

909

910 Figure 6 - Principal component analysis showing how the sites in Austria and Sweden differ  
911 based on the variables. PC1 and PC2 are the first two components, explaining most variance.  
912 The loadings (black arrows) are the weight of each variable in defining each principal  
913 component. The size of the arrows can tell how much they contribute defining this space, while  
914 the direction is their contribution on each axis. Tap: total annual precipitation; TxP: temperature  
915 x precipitation factor; pet: potential evapotranspiration; mat: mean annual temperature; tn: total  
916 soil nitrogen; soc: total soil organic carbon; cn: soil C:N ratio; ai: aridity index.